

From single-molecule fluorescence to photosynthesis with an STM

S. Cao¹, A. Roslawska¹, B. Doppagne¹, M. Roméo¹, M. Féron², F. Chérioux², H. Bulou¹, F. Scheurer¹,
G. Schull¹

¹ *Université de Strasbourg, CNRS, IPCMS, UMR 7504, Strasbourg, France*

² *Université Bourgogne Franche-Comté, FEMTO-ST, UFC, CNRS, Besançon, France*

Email: schull@unistra.fr

The electric current traversing the junction of a scanning tunneling microscope (STM) may lead to a local emission of light that can be used to generate sub-molecularly resolved fluorescence maps of individual molecules. Combined with spectral selection and time-correlated measurements, this hyper-resolved fluorescence microscopy approach allowed us to scrutinize the vibronic structure of individual molecules [1] in a very similar way than in the recent TERS reports, without requiring an optical excitation. We used this approach to characterize the photonics properties of charged species [2], to track the motion of hydrogen atoms within free-based phthalocyanine molecules [3], and more recently to follow resonance energy transfers between individual pigments, exploring processes occurring in photosynthetic complexes with sub-molecular spatial resolution [4].

These results constitute an important step towards photonic measurements with atoms-scale resolution [5].




Figure 1. Exploring energy transfers occurring in leaves with a scanning tunneling microscope.

- [1] B. Doppagne et al., *Phys. Rev. Lett.* 118, 127401 (2017)
- [2] B. Doppagne et al. *Science* 361, 251 (2018)
- [3] B. Doppagne et al. *Nature Nanotechnol.*15, 207 (2020).
- [4] S. Cao et al. *Nature Chem.* 12, 766 (2021)
- [5] A. Roslawska et al. *Physical Review X* 12, 011012 (2022)